

Abundance of Epiphytic Dinoflagellates from Coastal Waters off Jeju Island, Korea During Autumn 2009

Hyung Seop Kim¹, Wonho Yih^{2*}, Jong Hyeok Kim², Geumog Myung³, and Hae Jin Jeong⁴

¹Research Institute of Fisheries Technology, Kunsan 573-882, Korea

²Department of Oceanography, Kunsan National University, Kunsan 573-701, Korea

³Western Branch, Korea Fisheries Resources Agency, Kunsan 573-030, Korea

⁴School of Earth and Environmental Sciences, College of Natural Sciences, Seoul National University, Seoul 151-747, Korea

Received 23 June 2011; Revised 24 August 2011; Accepted 10 September 2011

© KSO, KORDI and Springer 2011

Abstract – The occurrence of harmful epiphytic dinoflagellates is of concern to scientists, the aquaculture industry, and government due to their toxicity not only to marine organisms but also to humans. There have been no studies on the abundance of the epiphytic dinoflagellates in Korean waters. We explored the presence of epiphytic dinoflagellates in the coastal waters off Jeju Island, southwestern Korea. Furthermore, we measured the abundance of epiphytic dinoflagellates on the thalli of 24 different macroalgae, collected from five different locations in October 2009. Five epiphytic dinoflagellate genera *Amphidinium*, *Coolia*, *Gambierdiscus*, *Ostreopsis*, and *Prorocentrum* were found. These five genera were observed on the thalli of the macroalgae *Chordaria flagelliformis*, *Martensia* sp., *Padina arborescens*, and *Sargassum* sp., while none were observed exceptionally on *Codium fragile*. The abundance of *Ostreopsis* spp. was highest on *Derbesia* sp. (8,660 cells/g wet weight), while that of *Gambierdiscus* spp. was highest on *Martensia* sp. (4,870 cells/g-ww). The maximum abundances of *Amphidinium* spp., *Coolia* spp., and *Prorocentrum* spp. were 410, 710, and 300 cells/g-ww, respectively. The maximum abundance of *Coolia* spp., *Gambierdiscus* spp., and *Ostreopsis* spp. obtained in the present study was lower than for other locations reported in literature. The results of the present study suggest that the presence and abundance of epiphytic dinoflagellates may be related to the macroalgal species of the coastal waters of Jeju Island.

Key words – *Amphidinium*, *Gambierdiscus*, HAB, *Ostreopsis*, *Prorocentrum*, red tide

1. Introduction

Species in the genera *Gambierdiscus*, *Ostreopsis*, *Coolia*, *Prorocentrum*, and *Amphidinium* are known to be epiphytic and/or benthic dinoflagellates (Taylor 1979, 1995, Hurbungs et al. 2001). Most of the epiphytic dinoflagellates is harmful to humans as well as to marine organisms, to which scientists, the aquaculture industry, and government are keenly attentive (Bagnis et al. 1980; Alcalá et al. 1988; Mangialajo et al. 2008). The majority of these dinoflagellates are known to be present in tropical or subtropical regions (Steidinger and Tangen 1997), but some species also live in the warm waters of temperate regions (Hurbungs et al. 2001; Pistocchi et al. 2011). The occurrence of epiphytic and benthic dinoflagellates in temperate waters has been reported as evidence of increasing water temperature (Graneli et al. 2011; Rhodes 2011).

Recently, Jeong et al. (submitted) reported the presence of *Coolia* spp. in the coastal waters of Jeju Island, southwestern Korea. However, this paper did not report the presence of the other epiphytic dinoflagellates at Jeju Island. We provide the first account of the occurrence of *Gambierdiscus*, *Ostreopsis* and epiphytic *Prorocentrum* and *Amphidinium* spp. from the coastal waters of Jeju Island. Additionally, we report the abundance of these epiphytic dinoflagellates, and the diverse macroalgal species on which they were found. The results of the present study should provide a basis for better understanding the ecology of epiphytic dinoflagellates in Korean waters.

*Corresponding author. E-mail: ywonho@kunsan.ac.kr

2. Materials and Methods

To measure the abundance of epiphytic dinoflagellates, macroalgal samples (24 different algal species) were collected by divers at 1–3 m depth off Hahnlím, Kosahn, Sahgye, Namwon, and Seongsan, Jeju Island, Korea (Fig. 1), on October 30 and November 1st, 2009. The water temperature and salinity were 21.0–23.6 °C and 29.2–31.6, respectively (Table 1). The samples were placed in plastic bags and then fixed with buffered 3% formaldehyde. Additional samples were placed in plastic bags, and then stored in ice-boxes, and then transported to the laboratory. The fixed macroalgal samples were vigorously shaken to detach dinoflagellates from the thalli of the algae and then screened through a 500 µm Nitex mesh. Filtrates were

concentrated to 1/5–1/10 after settling down the cells for 48 h. After being well mixed, all or >300 cells in one 1-ml Sedgwick-Rafter counting chamber were counted under a light microscope. The wet weight of each macroalgal sample was measured using a balance (Model CUW620H, CAS, Korea). Cells (not preserved) were collected from the macroalgal samples and photographed using a digital camera (Zeiss AxioCam HRc5, Carl Zeiss Ltd. Göttingen, Germany).

3. Results and Discussion

Five epiphytic dinoflagellate genera *Amphidinium*, *Coolia*, *Gambierdiscus*, *Ostreopsis*, and *Prorocentrum* were found in the coastal waters of Jeju island, Korea on October 31




Fig. 1. The five sampling sites at Jeju Island, southwestern Korea

Table 1. Water temperature (T, °C) and salinity (S) in the locations where macroalgae were collected on October 31 and November 1, 2009

Location	Latitude	Longitude	Temperature	Salinity
Hahnlím	N 33°24'25.46"	E 126°14'30.02"	21.1	28.9
Kosahn	N 33°17'52.30"	E 126°09'42.22"	21.8	31.1
Sahgye	N 33°13'34.24"	E 126°18'25.63"	23.6	31.6
Namwon	N 33°16'28.99"	E 126°42'56.96"	22.1	29.2
Seongsan	N 33°27'23.65"	E 126°55'45.37"	21.0	32.5




Fig. 2. Epiphytic dinoflagellates observed in this study. (A) *Amphidinium* sp. bar = 10 μ m. (B) *Gambierdiscus* sp. bar = 20 μ m. (C) *Ostreopsis* sp. bar = 20 μ m. (D) *Prorocentrum* sp. bar = 10 μ m

and November 1, 2009 (Fig. 2). All were observed on the thalli of the macroalgae *Chordaria flagelliformis*, *Martensia* sp., *Padina arborescens*, and *Sargassum* sp.. Furthermore, *Coolia*, *Gambierdiscus*, *Ostreopsis*, and *Prorocentrum* were observed on the thalli of the macroalgae *Ulva pertusa*, *Ecklonia caba*, and *Chondrus ocellatus*. However, none of these genera were observed with *Codium fragile* (Table 2). *Amphidinium* spp. were observed with seven macroalgal species (*Cladophora wrightiana*, *Sargassum* sp., *Dictyopteris divaricata*, *C. flagelliformis*, *P. arborescens*, and *Martensia* sp.), *Coolia* with seven macroalgal species (*U. pertusa*, *E. caba*, *Sargassum* sp., *C. flagelliformis*, *P. arborescens*, *Martensia* sp., and *C. ocellatus*), while *Prorocentrum* spp. with eight macroalgal species (*U. pertusa*, *E. caba*, *Sargassum* sp., *D. divaricata*, *C. flagelliformis*, *P. arborescens*, *Martensia* sp., *Gelidium amansii*, *Corallina* sp.). *Gambierdiscus* spp., were observed with all macroalgal species except *C. fragile* and *Sargassum siliquastrum*, while *Ostreopsis* spp. were observed with all macroalgal species except *C. fragile*,

Table 2. The abundance (cells/g wet weight) of the epiphytic dinoflagellates *Amphidinium* spp., *Coolia* spp., *Gambierdiscus* spp., *Ostreopsis* spp., *Prorocentrum* spp. on diverse macroalgae collected from 5 stations along the coasts of Jeju Island, Korea on October 31–November 1, 2009

	Macroalgae\Epiphytes	<i>Amphidinium</i>	<i>Coolia</i>	<i>Gambierdiscus</i>	<i>Ostreopsis</i>	<i>Prorocentrum</i>
Chlorophyta	<i>Cladophora wrightiana</i>	0–15	0	39	758	0
	<i>Ulva pertusa</i>	0	0–237	500	342	29–53
	<i>Cladophoropsis herpestica</i>	0	0	173	231	0
	<i>Derbesia</i> sp.	0	0	1,595	8,660	0
	<i>Codium fragile</i>	0	0	0	0	0
Phaeophyta	<i>Dictyopteris prolifera</i>	0	0	60	164	0
	<i>Ecklonia caba</i>	0	0–33	17–53	5–69	8
	<i>Dictyota okamurae</i>	0	0	1,000	185	0
	<i>Sargassum siliquastrum</i>	0	0	–	121	0
	<i>Sargassum</i> sp.	0–10	0–121	343	434	30
	<i>Dictyopteris divaricata</i>	0–37	0	13–200	87–216	12
	<i>Chordaria flagelliformis</i>	0–121	0–339	1,770	364	97
	<i>Zonaria diesingiana</i>	0	0	111	44–94	0
	<i>Padina arborescens</i>	0–137	0–652	789	583	171
	<i>Colpomenia sinuosa</i>	0–68	0	136	545	0
	<i>Dictyopteris undulata</i>	0	0	62	–	0
Rhodophyta	<i>Plocamium cartilagineum</i>	0	0	255	255	0
	<i>Pterocladia capillacea</i>	0	0	1,599	837	0
	<i>Lithothamnion</i> sp.	0	0	25	–	0
	<i>Martensia</i> sp.	0–406	0–710	4,871	3,349	304
	<i>Gelidium amansii</i>	0	0	229–426	78–688	233
	<i>Corallina</i> sp.	0	0	33–993	28–307	33
	<i>Chondrus ocellatus</i>	0	0–47	63	267	0
	<i>Plocamium telfairiae</i>	0	0	406–1477	50–966	0




Fig. 3. Mean abundance of epiphytic dinoflagellates in each of the five stations. Error bar represents standard error of mean

Dictyopteris undulata, and *Lithothamnion* sp. (Table 2). The data suggest that the presence of these five epiphytic dinoflagellate genera is affected by the macroalgal species in the study area.

The presence of the five epiphytic genera on the four macroalgal species, i.e. *C. flagelliformis*, *Martensia* sp., *P. arborescens*, and *Sargassum* sp., contrasts with their absence from a green alga, *C. fragile*. *Codium fragile*, however, was observed only at St. Kosahn where the coastline is more exposed than St. Namwon or Seongsan. Samples from St. Kosahn and Sahgye gave far lower abundance of epiphytic dinoflagellates than the other two

embayment stations (Fig. 3). The unexpected absence of epiphytic dinoflagellates from *Codium fragile* and the far lower abundance on macroalgae from more exposed sites may be related with species interactions of the epiphytes with physical and biological environment. Shaking effect due to turbulent water at the open coast (physical-biological interaction) and/or diverse bi-species interaction among different macroalga-epiphyte combinations may limit or support their success (Bomber et al. 1989).

The evidence suggests that *Derbesia* sp. is the optimal macroalgae for *Ostreopsis* spp., while *Martensia* sp. may be the optimal macroalgae for the other dinoflagellate genera studied herein (Table 2). However, abundance of epiphytic dinoflagellate estimated by cells per unit weight of substrate macroalgal blade tends to become higher on such macroalgal species with higher surface to volume ratio as *Derbesia* sp. and *Martensia* sp..

The maximum abundance of *Gambierdiscus* spp. ($\sim 5 \times 10^3$ cells/g ww) obtained in the present study is lower than that of *G. toxicus* from the Gambier Islands, Pacific Ocean (500×10^3 cells/g ww; Yasumoto et al. (1980)). The maximum abundance of *Ostreopsis* spp. ($\sim 9 \times 10^3$ cells/g ww) obtained in the present study is also quite lower than elsewhere (400 – $2,500 \times 10^3$ cells/g ww; Table 3). This might imply that the environmental conditions in the study area is at sub-optimal level to the five epiphytic dinoflagellate genera, which are presumably “settling down and adapting themselves” to the increasing water temperature of Jeju coastal waters.

Table 3. Comparison of the maximum abundances (MA, $\times 10^3$ cells/g wet weight) of the epiphytic dinoflagellates *Amphidinium* spp., *Coolia* spp., *Gambierdiscus* spp., *Ostreopsis* spp., *Prorocentrum* spp. obtained in the present study and those reported in literature

Genus	location	MA	Reference
<i>Amphidinium</i> spp.	Jeju Island, Korea	0.41	This study
<i>Coolia</i> spp.	Jeju Island, Korea	0.71	This study
<i>C. monotis</i>	Virgin Islands, Caribbean Sea	1,200	Calson & Tindall (1985)
<i>Gambierdiscus</i> spp.	Jeju Island, Korea	4.87	This study
<i>G. toxicus</i>	Gambier Islands, Pacific Ocean	500	Yasumoto et al.(1980)
<i>Ostreopsis</i> spp.	Jeju Island, Korea	8.66	This study
<i>O. ovata</i>	Genoa, Italy	2,541	Mangialajo et al. (2008)
<i>O. siamensis</i>	Auckland, New Zealand	1,406	Shears & Ross (2009)
<i>Ostreopsis</i> sp.	NW Mediterranean Sea	590	Vila et al. (2001)
<i>Ostreopsis</i> spp.	North Aegean Sea, Greece	405	Aligizaki & Nikolaidis (2006)
<i>Prorocentrum</i> spp.	Jeju Island, Korea	0.30	This study
<i>P. mexicanum</i>	Virgin Islands, Caribbean Sea	1,500	Calson & Tindall (1985)

Acknowledgements

We thank Mr. Yong-Deok Ko for technical supports in collecting samples from subsurface waters. This paper was supported by the Infrastructuring Grant for Marine Biotechnology Program funded by Korea Institute of Marine Science & Technology Promotion (KIMST)/Ministry of Land, Transportation and Marine Affairs (KMLTM) award to WH Yih, the National Research Foundation of Korea Grant funded by the Korea Government/MEST (NRF-C1ABA001-2010-0020700), and the Ecological Disturbance Research Program funded by KIMST/ KMLTM award to HJ Jeong.

References

- Alcala AC, Alcala LC, Garth JS, Yasumura D, Yasumoto T (1988) Human fatality due to ingestion of the crab *Demanda reynaudii* that contained a palytoxin-like toxin. *Toxicon* **26**:105-107
- Aligizaki K, Nikolaidis G (2006) The presence of the potentially toxic genera *Ostreopsis* and *Coolia* (Dinophyceae) in the North Aegean Sea, Greece. *Harmful Algae* **5**:717-730
- Bagnis R, Chanteau S, Chungue E, Hurtel JM, Yasumoto T, Inoue A (1980) Origins of ciguatera fish poisoning : A new dinoflagellate, *Gambierdiscus toxicus* Adachi and Fukuyo, definitively involved as a causal agent. *Toxicon* **18**:199-208
- Bomber JW, Rubio MG, Norris DR (1989) Epiphytism of dinoflagellates associated with the disease ciguatera: substrate specificity and nutrition. *Phycologia* **28**:360-368
- Carlson RD, Tindall DR (1985) Distribution and periodicity of toxic dinoflagellates in the Virgin Islands. In: Anderson DM, White AW, Baden DG (eds) *Toxic dinoflagellates*. Elsevier, New York, pp 171-176
- Granéli E, Vidyarthna NK, Funari E, Cumaranatunga P, Scenati R (2011) Can increases in temperature stimulate blooms of the toxic benthic dinoflagellate *Ostreopsis ovata*? *Harmful Algae* **10**:165-172
- Hurbungs MD, Jayabalan N, Chineah V (2001) Seasonal distribution of potentially toxic benthic dinoflagellates in the lagoon of Trou aux Biches, Mauritius. Food and Agricultural Research Council, Réduit, Mauritius, pp 211-217
- Koike K, Ishimaru T, Murano M (1991) Distributions of benthic dinoflagellates in Akajima Island, Okinawa, Japan. *Nippon Suisan Gakkaishi* **57**:2261-2264
- Mangialajo L, Bertolotto R, Cattaneo-Vietti C, Chiantore M, Grillo C, Lemee R, Melchiorre N, Moretto P, Povero P, Ruggieri N (2008) The toxic benthic dinoflagellate *Ostreopsis ovata*: Quantification of proliferation along the coastline of Genoa, Italy. *Mar Pollut Bull* **56**:1209-1214
- Pistocchi R, Pezzolesi L, Guerrini F, Vanucci S, Dell'Aversano C, Fattorusso E (2011) A review on the effects of environmental conditions on growth and toxin production of *Ostreopsis ovata*. *Toxicon* **57**:421-428
- Rhodes L (2011) World-wide occurrence of the toxic dinoflagellate genus *Ostreopsis*. Schmidt. *Toxicon* **57**:400-407
- Shears NT, Ross PM (2009) Blooms of benthic dinoflagellates of the genus *Ostreopsis* an increasing and ecologically important phenomenon on temperate reefs in New Zealand and worldwide. *Harmful Algae* **8**:916-925
- Steidinger KA, Tangen K (1997) Dinoflagellates. In: Tomas CR (ed) *Identifying marine phytoplankton*. Academic Press, Florida, USA pp 387-584
- Taylor FJR (1979) A description of the benthic dinoflagellate associated with maitotoxin and ciguatoxin, including observations on Hawaiian material. In: Taylor DL, Seliger HH (eds) *Toxic dinoflagellate blooms*. Elsevier North Holland, New York, pp 71-76
- Taylor FJR, Fukuyo Y, Larsen J (1995) Taxonomy of toxic dinoflagellates. In: Hallegraeff GM, Anderson DM, Cembella AD (eds) *Manual on harmful marine microalgae*. IOC manuals and guides, No.33, UNESCO, pp 283-317
- Vila M, Garcés E, Masó M (2001) Potentially toxic epiphytic dinoflagellate assemblages on macroalgae in the NW Mediterranean. *Aquat Microb Ecol* **26**:51-60
- Yasumoto T, Oshima Y, Murakami Y, Nagajima I, Bagnis R, Fukuyo Y (1980) Toxicity of benthic dinoflagellates found in coral reef. *Bull Japanese Soc Sci Fish* **46**:327-331